

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT AD-A216 547		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited.	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TK-89-1864	
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		7a. NAME OF MONITORING ORGANIZATION AFOSR	
6a. NAME OF PERFORMING ORGANIZATION West Virginia University	6b. OFFICE SYMBOL (if applicable)	7b. ADDRESS (City, State, and ZIP Code) Bldg 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Air Force Office of Scientific Research	8b. OFFICE SYMBOL (if applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER AFOSR-89-0068	
8c. ADDRESS (City, State, and ZIP Code) AFOSR/NM Building 410 Bolling AFB, DC 20332-6448		10. SOURCE OF FUNDING NUMBERS PROGRAM ELEMENT NO. PROJECT NO. TASK NO. WORK UNIT ACCESSION NO. AFOSR89-0068 2304 A8	
11. TITLE (Include Security Classification) Finding Critical Independent Sets and Critical Vertex Subsets are Polynomial Problems			
12. PERSONAL AUTHOR(S) Cun-Quan Zhang			
13a. TYPE OF REPORT Final Technical	13b. TIME COVERED FROM 11/88 TO 10/89	14. DATE OF REPORT (Year, Month, Day) 89/10/15	15. PAGE COUNT 15
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES FIELD GROUP SUB-GROUP		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number) Finding critical independent sets and critical vertex subsets are polynomial problems	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) An independent set J_c of a graph G is called critical if $ J_c - N(J_c) = \max \{ J - N(J) : J \text{ is an independent set of } G\}$ and a vertex subset U_c is called critical if $ U_c - N(U_c) = \max \{ U - N(U) : U \text{ is a subset of } V(G)\}$. In this paper, we will show that to find a critical independent set and a critical vertex subset of a graph are solvable in polynomial time.			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL DR. Neal Glassman		22b. TELEPHONE (Include Area Code) 22c. OFFICE SYMBOL (202) 767-5026 NM	

FINDING CRITICAL INDEPENDENT SETS
AND
CRITICAL VERTEX SUBSETS
ARE
POLYNOMIAL PROBLEMS

Cun-Quan Zhang*
Department of Mathematics
West Virginia University
Morgantown
West Virginia 26506 USA

ABSTRACT

An independent set J_c of a graph G is called critical if $|J_c| - |N(J_c)| = \max\{|J| - |N(J)| : J \text{ is an independent set of } G\}$, and a vertex subset U_c is called critical if $|U_c| - |N(U_c)| = \max\{|U| - |N(U)| : U \text{ is a vertex subset of } G\}$. In this paper, we will show that to find a critical independent set and a critical vertex subset of a graph are solvable in polynomial time.

Key words: Independent set, Polynomial algorithm
AMS subject classification: 05C35, 68R10
Abbreviated title: Critical independent sets

*. This research was partially supported by AFOSR under the grant 89-0068

S 1. INTRODUCTION

It has been proved by mathematicians that finding a maximum independent set in some certain kinds of graphs is solvable in polynomial time (for example, line graphs, bipartite graphs, circle graphs, circular arc graphs and claw free graphs, (see [GJ])) . But it is well-known that it is an NP-complete problem for general graphs, (see [GJS]) . In this paper, we will investigate another problem -- finding a certain kind of independent sets in general graphs. An independent set J_c of a graph G is called critical if $|J_c| - |N(J_c)|$ is the maximum of $|J| - |N(J)|$ over all independent sets J of G , where $N(J)$ is the set all vertices of G adjacent to some vertex of J . It will be proved in this paper that finding a critical independent set of a graph is solvable in polynomial time. Let

$$\alpha_c = \max \{ |J| - |N(J)| : J \text{ is an independent set of } G \}$$

which is a parameter of a graph G and is called the critical independence number of G . The critical independence number

α_c of a graph plays the central role in the study of fractional independence functions and fractional matching functions of graphs [GZ]. (It is proved in [GZ] that the fractional independence number and the fractional matching

number of a graph G are $\frac{n - \alpha_c}{2}$ and $\frac{n + \alpha_c}{2}$, respectively, where $n = |V(G)|$).

Some related problems and parameters of graphs will also be investigated in this paper. A vertex subset U_c of a graph $G = (V, E)$ is called critical if $|U_c| - |N(U_c)|$ is the maximum of $|U| - |N(U)|$ over all vertex subsets U of G . Let

$$\mu_c = \max \{ |U| - |N(U)| : U \text{ is a vertex subset of } G \}$$

which is a parameter of a graph G . Some similar parameters of graphs have been studied by Woodall [WD] and Mohar [MB].



For	<input checked="" type="checkbox"/>
ed	<input type="checkbox"/>
tion	<input type="checkbox"/>
lon/	
lity Codes	
avail and/or Special	

Dist
A-1

The binding number $b(G)$ [WD] and the isoperimetric number $i(G)$ [MB] of a graph G are defined as the following:

$$b(G) = \min \left\{ \frac{|N(U)|}{|U|} : U \subset V(G), U \neq \emptyset \text{ and } N(U) \neq V(G) \right\};$$

$$i(G) = \min \left\{ \frac{|\partial(U)|}{|U|} : U \subset V(G), U \neq \emptyset \text{ and } |U| \leq \frac{|V(G)|}{2} \right\}.$$

(where $\partial(U)$ is the number of edges of G with one endvertex in U). Later in this paper, it will be proved that $\alpha_c = \mu_c$.

Since the empty set is an independent set and the set of all vertices of a graph G is also a vertex subset of G , it is trivial that

$$\alpha_c \geq 0 \text{ and } \mu_c \geq 0$$

for any graph G . Note that the empty set and the entire graph are critical vertex subsets of some connected graph G if $\mu_c(G)=0$. If we are to avoid these two trivial vertex subsets \emptyset and $V(G)$, we may consider the following parameter of a graph G :

$$\mu'_c = \max \{ |U| - |N(U)| : U \subset V(G), U \neq \emptyset \text{ and } U \neq V(G) \}.$$

But for any connected graph G and a vertex v of G , we have that $N(V(G) \setminus \{v\}) = V(G)$ and therefore the parameter $\mu'_c(G)$ still have a lower bound -1 . In order to get more information about graphs, we prefer to only consider those vertex subsets U of a graph G such that $U \neq \emptyset$ and $N(U) \neq V(G)$ which is similar to the definition of the binding number of graphs. A vertex subset U of G is called proper if $U \neq \emptyset$ and $N(U) \neq V(G)$. A proper vertex subset U_{pc} of G is called critical if $|U_{pc}| - |N(U_{pc})|$ is the maximum of $|U| - |N(U)|$ over all proper vertex subsets U of G . The parameter μ_{pc} of a graph G is defined as the following:

$$\mu_{pc} = \max \{ |U| - |N(U)| : U \subset V(G), U \neq \emptyset \text{ and } U \neq V(G) \}$$

The problems will be proved to be solvable in polynomial time in this paper are listed as the following:

INSTANCE. let $G=(V,E)$ be a graph with the vertex set V and the edge set E and k be an integer.

PROBLEM 1. Is there an independent set J of G such that

$$|J| - |N(J)| \geq k?$$

PROBLEM 1*. Find a critical independent set J_c and the critical independence number α_c of G . (That is to find

$$\begin{aligned} \alpha_c &= |J_c| - |N(J_c)| \\ &= \max \{ |J| - |N(J)| : J \text{ is an independent set of } G \}. \end{aligned}$$

PROBLEM 2. Is there a vertex set U of G such that

$$|U| - |N(U)| \geq k?$$

PROBLEM 2*. Find a critical vertex subset U_c and the parameter μ_c of G . (That is to find

$$\begin{aligned} \mu_c &= |U_c| - |N(U_c)| \\ &= \max \{ |U| - |N(U)| : U \text{ is a vertex subset of } G \}. \end{aligned}$$

PROBLEM 3. Is there a proper vertex subset U of G such that

$$|U| - |N(U)| \geq k?$$

PROBLEM 3*. Find a critical proper vertex subset U_{pc} and the parameter μ_{pc} of G (That is to find

$$\begin{aligned} \mu_{pc} &= |U_{pc}| - |N(U_{pc})| \\ &= \max \{ |U| - |N(U)| : U \subset V(G), U \neq \emptyset \text{ and } U \neq V(G) \}. \end{aligned}$$

§ 2. MAIN RESULTS

THEOREM 1.

Problem 3 and 3* are solvable in polynomial time.

Before we prove the Theorem 1 we would like to consider the following problems first. And the Theorem 1 will be a corollary of the Theorem 2.

INSTANCE. Let $G=(V,E)$ be a graph with the vertex set V and the edge set E , $\{u,v\}$ be an ordered pair of nonadjacent vertices of G and k be an integer.

PROBLEM 4. Is there a vertex subset U of G such that
 $u \in U, v \notin N(U)$ and
 $|U| - |N(U)| \geq k$?

PROBLEM 4*. Find a vertex subset U_0 of G such that
 $|U_0| - |N(U_0)|$

$$= \max \{ |U| - |N(U)| : U \subset V(G), u \in U \text{ and } v \notin N(U) \}.$$

The vertex subset U_0 found in problem 4* is called (u,v) -critical subset of G .

THEOREM 2.

The problems 4 and 4* are solvable in polynomial time.

The following lemmas will be used in the proof of the Theorem 2.

LEMMA 3. (Hall's Theorem [HP])

Let $B=(V_1, V_2; E)$ be a bipartite graph. The graph B has a matching covering all vertices of V_2 if and only if
 $|U| \leq |N(U)|$ for any subset U of V_2 .

LEMMA 4.

Let $B = (V_1, V_2; E)$ be a bipartite graph. Assume that there is no matching of B covering all vertices of V_2 . We will have the following conclusions:

- (i). There is a subset U of V_2 such that $|N(U)| < |U|$;
- (ii). Let U_0 be a subset of V_2 such that $|U_0| - |N(U_0)|$ is as great as possible, then there is a matching of the induced bipartite subgraph $(U_0, N(U_0); E[U_0, N(U_0)])$ covering all vertices of $N(U_0)$.

Note, if A and B are a pair of disjoint vertex subset of a graph G, the set of all edges joining A and B is denoted by $E(A,B)$.

PROOF

The conclusion of (i) is an immediate corollary of Hall's Theorem.

Let U be a subset of V_2 such that $|U| - |N(U)|$ is as great as possible. Let $B'=(U, N(U); E[U, N(U)])$ be the subgraph of B induced by $U \cup N(U)$. We claim that $|X| \leq |N(X) \cap U|$ for any subset X of $N(U)$. If not, let $X \subseteq N(U)$ such that

$$|X| > |N(X) \cap U|.$$

We will consider the subset $Y = U \setminus N(X)$. Note that

$$N(Y) = N(U \setminus N(X)) \subseteq N(U) \setminus X.$$

And

$$\begin{aligned} |Y| - |N(Y)| &\geq |U \setminus N(X)| - |N(U) \setminus X| \\ &= [|U| - |U \cap N(X)|] - [|N(U)| - |X|] \\ &= [|U| - |N(U)|] + [|X| - |U \cap N(X)|] \\ &> |U| - |N(U)|. \end{aligned}$$

This contradicts the choice of U that $|U| - |N(U)|$ is maximum. So by Hall's theorem, there is a matching in B' which covers all vertices of $N(U)$.

###

PROOF OF THE THEOREM 2.

We are only to prove that problem 4* is solvable in polynomial time. Let $G=(V,E)$ be a graph with the vertex set $V=\{1,2,3,\dots,n\}$ and the edge set E. We will consider the ordered pair of vertices (1,2) of G and find a (1,2)-critical vertex subset.

Define a bipartite graph $B = (X,Y; E_B)$ where

$$X = \{x_1, \dots, x_n\},$$

$$Y = \{y_1, \dots, y_n\}$$

and

$$E_B = \{(x_i, y_j) : (i, j) \text{ is an edge of } G\}.$$

Let V' be a subset of $V(G)$, then the corresponding subsets in X and Y are denoted by $X(V')$ and $Y(V')$, respectively. For example, if $V' = \{i_1, \dots, i_t\}$, then

$$X(V') = X(\{i_1, \dots, i_t\}) = \{x_{i_1}, \dots, x_{i_t}\} \text{ and}$$

$$Y(V') = Y(\{i_1, \dots, i_t\}) = \{y_{i_1}, \dots, y_{i_t}\}. \quad (\text{Here } X \text{ and } Y \text{ can be}$$

considered as bijections mapping $\{1, 2, \dots, n\}$ onto $\{x_1, \dots, x_n\}$

and $\{y_1, \dots, y_n\}$). If $W = \{x_{i_1}, \dots, x_{i_t}\} \subset X$ (or

$W = \{y_{i_1}, \dots, y_{i_t}\} \subset Y$), then the corresponding subset $\{i_1, \dots, i_t\}$

of $V(G)$ is denoted by $X^{-1}(W)$ (or $Y^{-1}(W)$, respectively). The

set of all neighbors of a vertex u in B is denoted by $N_B(u)$.

If i is a vertex of G , then

$$N_B(x_i) = \{y_j \in Y : (x_i, y_j) \in E_B\} = \{y_j \in Y : (i, j) \in E(G)\} = Y(N(i)).$$

A weight $w: X \cup Y \rightarrow [0, 2]$ is called a (1,2)-proper weight of B if

$$w(x_1) = 2,$$

$$w(y_2) = 1,$$

$$1 \leq w(x_i) \leq 2 \quad \text{for each vertex } x_i \in X,$$

$$0 \leq w(y_i) \leq 1 \quad \text{for each vertex } y_i \in Y$$

$$\text{and} \quad 0 \leq w(x_i) + w(y_i) \leq 2 \quad \text{for each edge } (x_i, y_i) \in E_B.$$

The total weight $\sum_{u \in X \cup Y} w(u)$ of B is denoted by $w(B)$. A (1,2)-

proper weight w_m of B is call optimum if

$$w_m(B) = \max\{w(B) : w \text{ is a (1,2)-proper weight of } B\}.$$

It is obvious that finding an optimum (1,2)-proper weight of B is a linear programming problem. Hence it is solvable in polynomial time. The purpose of the investigation of an optimum (1,2)-proper weight w_m of B is to prove that the vertex subset $\{i \in V(G) : w_m(x_i) > 1\}$ is a (1,2)-critical subset of G .

I. Let w_m be an optimum $(1,2)$ -proper weight of B and U_0 be a $(1,2)$ -critical vertex subset of G . We claim that

$$w_m(B) \geq 2n + |U_0| - |N(U_0)| = 2n + \alpha_c \quad (1)$$

Consider the following weight w_1 of B :

$$w_1(x_i) = \begin{cases} 2 & \text{if } i \in U_0 \\ 1 & \text{otherwise} \end{cases}$$

$$\text{and } w_1(y_j) = \begin{cases} 0 & \text{if } j \in N(U_0) \\ 1 & \text{otherwise} \end{cases}.$$

It is easy to see that w_1 is a $(1,2)$ -proper weight of B and

$$w_1(B) = 2|U_0| + (n - |U_0|) + (n - |N(U_0)|) \\ = 2n + |U_0| - |N(U_0)|.$$

By the choice of w_m , we have verified the inequality (1).

II. Let

$$X_b = \{x_i : w_m(x_i) > 1\}$$

$$Y_s = \{y_i : w_m(y_i) < 1\}$$

$$X_{b'} = \{x_i : 1 < w_m(x_i) < 2\}$$

and

$$Y_{s'} = \{y_i : 0 < w_m(y_i) < 1\}.$$

By the definition of $(1,2)$ -proper weight, it is obvious that $N_B(X_b) \subseteq Y_s$ and $N_B(Y_{s'}) \subseteq [X \setminus X_b] \cup X_{b'}$.

III. Case 1. Suppose that there is a matching M in the induced subgraph $B(X_{b'} \cup Y_{s'})$ covering all vertices of $Y_{s'}$. We claim that X_b is a $(1,2)$ -critical vertex subset in this case. We are to adjust the weight w_m so that the new weight of each vertex in $X_{b'}$ is two and the new weight of each vertex in $Y_{s'}$ is one. If we can verify that this new weight is optimum, by the inequality (1), it can be shown that $X^{-1}(X_b)$ is a $(1,2)$ -critical vertex set of G .

If (u,v) is an edge of M , then let $u=M(v)$ and $v=M(u)$. And the sets of vertices of $X_{b'}$ covered and not covered by M is denoted by $M(Y_{s'})$ and $X_{b'} \setminus M$, respectively. Thus $X_{b'} \cup Y_{s'} = Y_{s'} \cup M(Y_{s'}) \cup (X_{b'} \setminus M)$ since M covers all vertices of $Y_{s'}$.

Consider the following weight w_2 of B :

$$w_2(v) = \begin{cases} 2 & \text{if } v \in X_b, \\ 0 & \text{if } v \in Y_s, \\ w_m(v) & \text{otherwise} \end{cases}.$$

Since any vertex of Y adjacent to a vertex x_i of X_b must be in Y_s in which the weight of each vertex is zero, w_2 is a $(1,2)$ -proper weight of B . Note that w_m is optimum and the total weight $w_m(B)$ cannot be less than $w_2(B)$. We claim that w_2 is also an optimum $(1,2)$ -proper weight of B by proving that $w_m(B) \leq w_2(B)$. Since

$$\{v \in X \cup Y: w_m(v) \neq w_2(v)\} = X_b \cup Y_s,$$

we must have that

$$\begin{aligned} w_m(B) - w_2(B) &= \sum_{v \in X_b} [w_m(v) - w_2(v)] + \sum_{v \in Y_s} [w_m(v) - w_2(v)] \\ &= \sum_{v \in Y_s} \{ [w_m(v) - w_2(v)] + [w_m(M(v)) - w_2(M(v))] \} + \\ &\quad + \sum_{v \in X_b \setminus M} [w_m(v) - w_2(v)]. \end{aligned}$$

$$\begin{aligned} \text{Here} \quad & [w_m(v) - w_2(v)] + [w_m(M(v)) - w_2(M(v))] \\ &= [w_m(v) - w_m(M(v))] - [w_2(v) - w_2(M(v))] \\ &\leq 2 - (0+2) \\ &= 0 \end{aligned}$$

for any $v \in Y_s$, and

$$w_m(v) - w_2(v) < 2 - 2 = 0$$

for any $v \in X_b \setminus M$. This implies that $w_m(B) - w_2(B) \leq 0$.

Therefore w_2 is also an optimum $(1,2)$ -proper weight of B and $w_m(B) = w_2(B)$. The total weight of w_2 is

$$w_2(B) = 2|X_b| + |X \setminus X_b| + |Y \setminus Y_s|.$$

Since $N_B(X_b) \subseteq Y_s$, we must have that

$$\begin{aligned} w_m(B) = w_2(B) &\leq 2|X_b| + (n - |X_b|) + (n - |N(X_b)|) \\ &= 2n + |X_b| - |N(X_b)| \\ &= 2n + |X^{-1}(X_b)| - |N(X^{-1}(X_b))| \\ &\leq 2n + \alpha_c \quad (\text{by the definition} \end{aligned}$$

of α_c).

By (1), all equalities hold and therefore $X^{-1}(X_b) = \{i \in V(G) : w_m(x_i) > 1\}$ is a (1,2)-critical vertex subset of G .

IV. Case 2. If there is no matching of $B(X_b' \cup Y_{s'})$ covering all vertices of $Y_{s'}$. By Lemma 4, there is a subset Y_0 of $Y_{s'}$ such that (i). $|Y_0| > |N_B(Y_0) \cap X_b'|$ and, (ii). there is a matching M' in the induced bipartite subgraph $B(Y_0 \cup [N_B(Y_0) \cap X_b'])$ covering all vertices of $N_B(Y_0) \cap X_b'$. We are to adjust the weight w_m so that the new weight of each vertex in $Y_0 \cup [N_B(Y_0) \cap X_b']$ is one. We will find that the new weight is greater than w_m . It will contradict that w_m is optimum.

Consider the following weight w_3 :

$$w_3(v) = \begin{cases} 1 & \text{if } v \in Y_0 \cup [N_B(Y_0) \cap X_b'] \\ w_m(v) & \text{otherwise} \end{cases}.$$

The weight w_3 is (1,2)-proper since $x_1 \notin N(Y_{s'})$ and any vertex adjacent to a vertex of Y_0 must be in $[X \setminus X_b] \cup [N_B(Y_0) \cap X_b']$ in which the weight of each vertex is one. Note that w_m is an optimum (1,2)-proper weight of B , thus the total weight of w_m cannot be less than the total weight of w_3 . Since

$$\begin{aligned} \{v \in V(B) : w_m(v) \neq w_3(v)\} &\subseteq Y_0 \cup [N_B(Y_0) \cap X_b'] \\ &= [N_B(Y_0) \cap X_b'] \cup M'[N_B(Y_0) \cap X_b'] \cup [Y_0 \setminus M'], \end{aligned}$$

where $M'[N_B(Y_0) \cap X_b']$ and $[Y_0 \setminus M']$ are the sets of vertices of Y_0 covered and not covered by M' , respectively. Thus we must have that

$$\begin{aligned} w_m(B) - w_3(B) &= \sum_{v \in N_B(Y_0) \cap X_b'} [w_m(v) - w_3(v)] + \sum_{v \in Y_0} [w_m(v) - w_3(v)] \\ &= \sum_{v \in N_B(Y_0) \cap X_b'} \{ [w_m(v) - w_3(v)] + [w_m(M(v)) - w_3(M(v))] \} + \\ &\quad + \sum_{v \in Y_0 \setminus M'} [w_m(v) - w_3(v)]. \end{aligned}$$

But

$$[w_m(v) - w_3(v)] + [w_m(M(v)) - w_3(M(v))]$$

$$\begin{aligned}
&= [w_m(v) - w_m(M(v))] - [w_3(v) - w_3(M(v))] \\
&\leq 2 - (1+1) \\
&= 0
\end{aligned}$$

for any $v \in N_B(Y_0) \cap X_{b'}$, and

$$w_m(v) - w_3(v) < 1-1 = 0$$

for any $v \in Y_0 \setminus M'$. Since $|Y_0| > |N_B(Y_0) \cap X_{b'}|$, the set $Y_0 \setminus M'$ is not empty and we have that $w_m(B) - w_3(B) < 0$. This contradicts that w_m is optimum and completes the proof of the theorem.

###

Proof of the Theorem 1.

For any ordered pair of non-adjacent vertices $\{u, v\}$ of G , by Theorem 2, we can find a (u, v) -critical subset in polynomial time. The (u, v) -critical subset is denoted by $U_C(u, v)$. Then choose a (u_0, v_0) -critical vertex subset $U_C(u_0, v_0)$ such that

$$\begin{aligned}
&|U_C(u_0, v_0)| - |N(U_C(u_0, v_0))| \\
&= \max\{|U_C(u, v)| - |N(U_C(u, v))| : (u, v) \text{ are an ordered}
\end{aligned}$$

pair of nonadjacent vertices of $G\}$ which is a proper critical subset of G desired in problem 3*. The total cost of finding a critical proper vertex subset is polynomial since the cost of finding a (u, v) -critical vertex subset is polynomial and the number of pairs of non-adjacent vertices in G is at most $\binom{n}{2}$.

###

Theorem 5.

Problem 2 and 2* are solvable in polynomial time.

Proof.

Let $G=(V, E)$ be a graph. Consider a new graph G' by adding two isolated vertices x, y to G . Let U_C be a (x, y) -

critical subset of G' . Obviously, $x, y \in U_c$ and it is clear that $U_c \setminus \{x, y\}$ is a critical subset of G .

Alternating proof of the Theorem: Define a bipartite graph $B = (X, Y; E_B)$ where $X = \{x_1, \dots, x_n\}$, $Y = \{y_1, \dots, y_n\}$ and $E_B = \{(x_i, y_j) : (i, j) \text{ is an edge of } G\}$. And assign a weight w to the vertex set of B such that $w: X \cup Y \rightarrow [0, 2]$ and

$$\begin{aligned} 1 \leq w(x_i) \leq 2 & \quad \text{for each vertex } x_i \in X, \\ 0 \leq w(y_i) \leq 1 & \quad \text{for each vertex } y_i \in Y \\ \text{and} \quad 0 \leq w(x_i) + w(y_i) \leq 2 & \quad \text{for each edge } (x_i, y_i) \in E_B. \end{aligned}$$

The total weight $\sum_{u \in X \cup Y} w(u)$ is denoted by $w(B)$. Let

$$w_m(B) = \max\{w(B) : w \text{ is a weight of } B \text{ satisfies the above definition}\}.$$

By an argument similar to the proof of Theorem 2, we can prove that the set $V_b = \{i \in V(G) : w_m(x_i) > 1\}$ is an critical vertex subset of G .

###

Theorem 6.

Problem 1 and 1^* are solvable in polynomial time.

Before the proof of the Theorem 6, we will prove a Theorem by which Theorem 6 is an immediate corollary.

Theorem 7.

Let $G = (V, E)$ be a graph.

(i). Let U_c be a critical vertex subset of G and T_1, \dots, T_h be all non-trivial components of the induced subgraph $G(U_c)$. Then $J = V(U_c) \setminus [V(T_1) \cup \dots \cup V(T_h)]$ is a critical independent set of G and $|J| - |N(J)| = |U_c| - |N(U_c)|$.

(ii). $\max \{ |J| - |N(J)| : J \text{ is an independent set of } G \}$
 $= \max \{ |U| - |N(U)| : U \text{ is a vertex subset of } G \}$.

That is, any critical independent set is also a critical vertex subset of G and therefore $\alpha_c = \mu_c$.

Proof.

It is obvious that

$$0 \leq \alpha_c = \max \{ |J| - |N(J)| : J \text{ is an independent set of } G \}$$

$$\leq \max \{ |U| - |N(U)| : U \text{ is a vertex subset of } G \} = \mu_c$$

.....(2)

since any independent set is a vertex subset of G .

Let U_c be a critical vertex subset of G . The Theorem is trivial if U_c is an empty set. Assume that U_c is a counterexample to the Theorem containing minimum number of vertices. By the assumption U_c cannot be an independent set of G . Let T be a non-trivial component of the induced subgraph $G(U_c)$. It is clear that $V(T) \subseteq N(T)$ since T is not a singleton. Thus

$$|U_c \setminus T| - |N(U_c \setminus T)| \geq [|U_c| - |T|] - [|N(U_c)| - |T|]$$

$$= |U_c| - |N(U_c)|$$

It implies that $U_c \setminus T$ is also a critical vertex subset of G which contains less number of vertices than U_c . It contradicts the choice of U_c and therefore completes the proof of the Theorem.

###

Proof of the Theorem 6.

Let U_c be a critical vertex subset of G . Let T_1, \dots, T_t be all non-trivial components of the induced subgraph $G(U_c)$. Then by Theorem 7, $U_c \setminus \{T_1, \dots, T_t\}$ is a critical independent

set of G . Since finding a critical vertex subset U_c and deleting all vertices of U_c incident with some edge of $G(U_c)$ only need polynomial cost, finding a critical independent set is solvable in polynomial time.

Alternating proof of the Theorem (also see [GZ]):

Consider a weight $w: V(G) \rightarrow [0,1]$ such that

$0 \leq w(v) \leq 1$ for each vertex v of G
 and $w(u) + w(v) \leq 1$ for each edge (u,v) of G

The total weight $\sum_{u \in V(G)} w(u)$ is denoted by $w(G)$. Let

$w_m(G) = \max\{w(G) : w \text{ is a weight of } G \text{ satisfies the above definition}\}.$

Obviously finding w_m is a linear programming problem. By an argument similar to the proof of Theorem 2, we can prove that the set $V_b = \left\{ v \in V(G) : w_m(v) > \frac{1}{2} \right\}$ is an critical independent set of G .

(Note that the weight w defined above is called a fractional independence function of a graph G which was introduced by Domke, Hedetniemi, Laskar in [DHL] and by Grinstead, Slater in [GS2], and were studied by Grinstead, Slater in [GS1] and by Zhang in [CG].)

###

Note, S. Poljak (personal communication) recently suggested an alternative proof of Theorem 2 by the theory of integer programming.

REFERENCES

- DHL Domke, G.S., Hedetniemi, S.T. and Laskar, R.C., Fractional Packings, Coverings, and Irredundance in graphs, preprint.
- GJ Garey, M.R. and Johnson, D.S., Computers and intractability, W.H. Freeman and Company, New York 1979

- GJS Garey, M.R., Johnson, D.S. and Stockmeyer, L. Some simplified NP-Complete graph problems, Theor. Comput. Sci. **1** (1976) 237-267
- GS1 Grinstead, D.L. and Slater, P.J., On fractional covering and fractional independence parameters, Third Carbondale Combinatorics Conference, (1988)
- GS2 Grinstead, D.L. and Slater, P.J., Fractional domination and fractional packings in graphs, preprint.
- GZ Goldwasser, J. and Zhang, C.Q., Fractional independence number and fractional matching number, preprint.
- HP Hall, P., On representatives of subsets. J. London math. Soc., **10**, (1935) 26-30.
- MB Mohar, B., Isoperimetric number of graphs, preprint.
- MG Minty, G.J., On maximal independent sets of vertices in claw-free graphs. J. Combinatorial Theory, B **28** (1977) 283-304
- WD Woodall, D.R., The binding number of a graph and its Anderson number. J. Combinatorial Theory (B), **15**, 225-255.